



<http://www.womenshealth.gov>

1-800-994-9662

TDD: 1-888-220-5446

Menstruation and the Menstrual Cycle

Q: What is menstruation?

A: Menstruation (men-STRAY-shuhn) is a woman's monthly bleeding. When you menstruate, your body sheds the lining of the uterus (womb). Menstrual blood flows from the uterus through the small opening in the cervix and passes out of the body through the vagina (see diagram on page 2). Most menstrual periods last from 3 to 5 days.

Q: What is the menstrual cycle?

A: When periods (menstruations) come regularly, this is called the menstrual cycle. Having regular menstrual cycles is a sign that important parts of your body are working normally. The menstrual cycle provides important body chemicals, called hormones, to keep you healthy. It also prepares your body for pregnancy each month. A cycle is counted from the first day of 1 period to the first day of the next period. The average menstrual cycle is 28 days long. Cycles can range anywhere from 21 to 35 days in adults and from 21 to 45 days in young teens.

The rise and fall of levels of hormones during the month control the menstrual cycle.

Q: What happens during the menstrual cycle?

A: In the first half of the cycle, levels of estrogen (the “female hormone”) start to rise. Estrogen plays an important role in keeping you healthy, especially by helping you to build strong bones and to help keep them strong as you get older. Estrogen also makes the lining of the uterus (womb) grow and thicken. This lining of the womb is a place that will nourish the embryo if a pregnancy occurs. At the same time the lining of the womb is growing, an egg, or ovum, in one of the ovaries starts to mature. At about day 14 of an average 28-day cycle, the egg leaves the ovary. This is called ovulation.

After the egg has left the ovary, it travels through the fallopian tube to the uterus. Hormone levels rise and help prepare the uterine lining for pregnancy. A woman is most likely to get pregnant during the 3 days before or on the day of ovulation. Keep in mind, women with cycles that are shorter or longer than average may ovulate before or after day 14.

A woman becomes pregnant if the egg is fertilized by a man’s sperm cell and attaches to the uterine wall. If the egg is not fertilized, it will break apart. Then, hormone levels drop, and the thickened lining of the uterus is shed during the menstrual period.




<http://www.womenshealth.gov>

1-800-994-9662

TDD: 1-888-220-5446

The picture below shows the path the egg takes from the ovary, through the fallopian tube, and to the uterus.



Q: What is a typical menstrual period like?

A: During your period, you shed the thickened uterine lining and extra blood through the vagina. Your period may not be the same every month. It may also be different than other women's periods. Periods can be light, moderate, or heavy in terms of how much blood comes out of the vagina. This is called menstrual flow. The length of the period also varies. Most periods last from 3 to 5 days. But, anywhere from 2 to 7 days is normal.

For the first few years after menstruation begins, longer cycles are common. A woman's cycle tends to shorten and become more regular with age. Most of the time, periods will be in the range of 21 to 35 days apart.

Q: What kinds of problems do women have with their periods?

A: Women can have a range of problems with their periods, including pain, heavy bleeding, and skipped periods.

- **Amenorrhea (ay-men-uh-REE-uh)** — the lack of a menstrual period. This term is used to describe the absence of a period in:
 - Young women who haven't started menstruating by age 15
 - Women and girls who haven't had a period for 90 days, even if they haven't been menstruating for long

Causes can include:

- Pregnancy
- Breastfeeding
- Extreme weight loss
- Eating disorders
- Excessive exercising
- Stress
- Serious medical conditions in need of treatment

As above, when your menstrual cycles come regularly, this means that important parts of your body are working normally. In some cases, not having menstrual periods can mean that your ovaries have stopped producing normal amounts of estrogen. Missing these hormones can have important effects on your overall health. Hormonal problems, such as those caused by polycystic ovarian syndrome (PCOS) or serious problems with the reproductive organs, may be involved. It's important to talk to a doctor if you have this problem.



<http://www.womenshealth.gov>

1-800-994-9662

TDD: 1-888-220-5446

- **Dysmenorrhea (dis-men-uh-REE-uh)** — painful periods, including severe cramps. Menstrual cramps in teens are caused by too much of a chemical called prostaglandin (pros-tuh-GLAN-duhn). Most teens with dysmenorrhea do not have a serious disease, even though the cramps can be severe. In older women, the pain is sometimes caused by a disease or condition such as uterine fibroids or endometriosis.

For some women, using a heating pad or taking a warm bath helps ease their cramps. Some over-the-counter pain medicines can also help with these symptoms. They include:

- Ibuprofen (eye-byu-PROH-fuhn) (for instance, Advil, Motrin, Midol Cramp)
- Ketoprofen (key-toh-PROH-fuhn) (for instance, Orudis KT)
- Naproxen (nuh-PROK-suhn) (for instance, Aleve)

If these medicines don't relieve your pain or the pain interferes with work or school, you should see a doctor. Treatment depends on what's causing the problem and how severe it is.

- **Abnormal uterine bleeding** — vaginal bleeding that's different from normal menstrual periods. It includes:
 - Bleeding between periods
 - Bleeding after sex
 - Spotting anytime in the menstrual cycle
 - Bleeding heavier or for more days than normal
 - Bleeding after menopause

Abnormal bleeding can have many causes. Your doctor may start by checking for problems that are most common in your age group. Some of them are not serious and are easy to treat. Others can be more serious. Treatment for abnormal bleeding depends on the cause.

In both teens and women nearing menopause, hormonal changes can cause long periods along with irregular cycles. Even if the cause is hormonal changes, you may be able to get treatment. You should keep in mind that these changes can occur with other serious health problems, such as uterine fibroids, polyps, or even cancer. See your doctor if you have any abnormal bleeding.

Q: When does a girl usually get her first period?

A: In the United States, the average age for a girl to get her first period is 12. This does not mean that all girls start at the same age. A girl can start her period anytime between the ages of 8 and 15. Most of the time, the first period starts about 2 years after breasts first start to develop. If a girl has not had her first period by age 15, or if it has been more than 2 to 3 years since breast growth started, she should see a doctor.

For help teaching your daughter about her changing body, visit <http://www.girlshealth.gov/body/period/index.cfm> for information on getting your period.

Q: How long does a woman have periods?

A: Women usually have periods until menopause. Menopause occurs between the ages of 45 and 55, usually around age 50. Menopause means that a woman is no longer ovulating (produc-



<http://www.womenshealth.gov>

1-800-994-9662

TDD: 1-888-220-5446

ing eggs) or having periods and can no longer get pregnant. Like menstruation, menopause can vary from woman to woman and these changes may occur over several years.

The time when your body begins its move into menopause is called the menopausal transition. This can last anywhere from 2 to 8 years. Some women have early menopause because of surgery or other treatment, illness, or other reasons. If you don't have a period for 90 days, you should see your doctor. He or she will check for pregnancy, early menopause, or other health problems that can cause periods to stop or become irregular.

Q: When should I see a doctor about my period?

A: See your doctor about your period if:

- You have not started menstruating by the age of 15.
- You have not started menstruating within 3 years after breast growth began, or if breasts haven't started to grow by age 13.
- Your period suddenly stops for more than 90 days.
- Your periods become very irregular after having had regular, monthly cycles.
- Your period occurs more often than every 21 days or less often than every 35 days.
- You are bleeding for more than 7 days.
- You are bleeding more heavily than usual or using more than 1 pad or tampon every 1 to 2 hours.
- You bleed between periods.
- You have severe pain during your period.

- You suddenly get a fever and feel sick after using tampons.

Q: How often should I change my pad and/or tampon?

A: You should change a pad before it becomes soaked with blood. Each woman decides for herself what works best. You should change a tampon at least every 4 to 8 hours. Make sure to use the lowest absorbency tampon needed for your flow. For example, use junior or regular tampons on the lightest day of your period. Using a super absorbency tampon on your lightest days increases your risk for toxic shock syndrome (TSS). TSS is a rare but sometimes deadly disease. TSS is caused by bacteria that can produce toxins. If your body can't fight the toxins, your immune (body defense) system reacts and causes the symptoms of TSS (see below).

Young women may be more likely to get TSS. Using any kind of tampon puts you at greater risk for TSS than using pads. The Food and Drug Administration (FDA) recommends the following tips to help avoid tampon problems:

- Follow package directions for insertion.
- Choose the lowest absorbency for your flow.
- Change your tampon at least every 4 to 8 hours.
- Consider switching between pads and tampons.
- Know the warning signs of TSS (see below).
- Don't use tampons between periods.



<http://www.womenshealth.gov>

1-800-994-9662

TDD: 1-888-220-5446

If you have any of these symptoms of TSS while using tampons, take the tampon out, and contact your doctor right away:

- Sudden high fever (over 102 degrees)
- Muscle aches
- Diarrhea

- Vomiting
- Dizziness and/or fainting
- Sunburn-like rash
- Sore throat
- Bloodshot eyes ■

For more information

For more information about menstruation, contact [womenshealth.gov](http://www.womenshealth.gov) at 1-800-994-9662 or the following organizations:

Food and Drug Administration

Phone Number(s): (888) 463-6332

Internet Address: www.fda.gov

American College of Obstetricians and Gynecologists (ACOG) Resource Center

Phone Number(s): (202) 638-5577; (202) 863-2518 (for publications requests only)

Internet Address: www.acog.org

Association of Reproductive Health Professionals

Phone Number(s): (202) 466-3825

Internet Address: <http://www.arhp.org>

Center for Young Women's Health

Phone Number(s): (617) 355-2994

Internet Address: <http://www.youngwomenshealth.org>

All material contained in this FAQ is free of copyright restrictions, and may be copied, reproduced, or duplicated without permission of the Office on Women's Health in the Department of Health and Human Services. Citation of the source is appreciated.

This FAQ was reviewed by:

Lawrence M. Nelson, M.D.

Investigator

National Institute of Child Health and Human Development

National Institutes of Health

Content last updated October 21, 2009.